Incentivized Kidney Exchange

Tayfun Sonmez[y] M. Utku Unver[z] M. Bumin Yenmez[x]

1 Introduction

While transplantation is the best remedy for end-stage renal disease, a severe shortage of transplant kidneys persists worldwide. Kidneys for transplantation can be harvested from deceased or living donors. As of February 2018, more than 280,000 kidney transplants from deceased donors and more than 145,000 from living donors have been performed in the US. The number of willing living donors has been considerably higher than the number of living-donor transplants performed, but intended gifts of a large fraction of potential donors have not materialized due to biological incompatibilities. More than thirty percent of potential living donors are blood-type incompatible, and about five percent are tissue-type incompatible, with their intended recipients. Blood type O patients are especially disadvantaged by these biological barriers because they are blood-type compatible with type O donors only. In contrast, blood type A patients are blood-type compatible with donors of types A and O, blood type B patients are blood-type compatible with donors of types B and O, and blood type AB patients are blood-type compatible with donors of all blood types.[1] The resulting disadvantage to type O patients is mitigated in deceased-donor transplants by a policy that reserves type O kidneys for type O patients, but a similar policy is not possible for living-donor transplants since a living-donor kidney is intended as a gift for a loved one.

Kidney exchange emerged as a transplantation modality over the last 15 years to lessen the prohibitive effects of biological barriers on living-donor transplantation. In its most basic form, a kidney exchange is a swap of donors between two patients who are each incompatible with their own donor but compatible with the other patient's donor. Both donors' intended gifts are materialized through the exchange, providing each patient with a transplant. However, type O patients are again less likely to benefit from this transplantation modality. Consider a type O patient unable to receive a transplant from his blood-type incompatible type A donor. The pair can potentially swap donors with a type A patient who has a type O donor; but being blood-type compatible, these pairs are rare, only arriving when they are tissue-type incompatible. Hence, a large number of "underdemanded" type O patients with type A donors compete for a relatively scarce population of "overdemanded" type A patients with type O donors.[2] Ironically, these pairs with highly sought-after type O donors become available for exchange only because of a tissue-type incompatibility. A biological barrier to transplantation results in an increase in the number of living-donor transplants by facilitating a welfare-increasing utilization of living donors.

Of course the competition for an exchange would not be so unfavorable for type O patients with blood-type incompatible donors if all pairs participated in kidney exchange rather than only incompatible pairs. Indeed, when a clearinghouse for organized kidney exchange was initially proposed, market designers advocated a mechanism where all pairs would participate in exchange, whether they are compatible or not (Roth, Sönmez, and Ünver, 2004). However, since patients with compat-

[1] For the US, 45.6 percent of the population is blood type O, 37.8 percent of the population is blood type A, 12.6 percent of the population is blood type B, and 4 percent of the population is blood type AB.

[2] Based on 2012-2014 data from the three largest kidney-exchange clearinghouses in the US, the percentage of pairs with type O patients was in the range 58.4–60.7 percent and the percentage of pairs with type O donors was in the range 30.8–33 percent (Agarwal et al., 2017).

ible donors are in no need of an exchange for a transplant, the practice of kidney exchange evolved

equity in access for living-donor transplantation, mainly by increasing transplants to type O patients through donor exchanges with incentivized pairs. This in turn also improves equity in access for deceased-donor transplantation, since type O patients who bene t from incentivized exchange no longer compete for deceased-donor transplants. Blood types O and B are more heavily represented among minority groups, and hence any disadvantage to patients of these blood types also correlates with an inequity of access across di erent ethnic backgrounds. As such, incentivized exchange also reduces disparities across racial and ethnic groups. To our knowledge, our proposed policy is the rst to enhance both the e ciency and equity of the system.[6]

To analyze the e ciency and equity implications of incentivized exchange, we introduce a new and analytically tractable dynamic large-market model of kidney transplantation[7]. Unlike former models that focus on a single organ-allocation technology, our model can be used to analyze the

literature also explored providing incentives through exchanging the donor of a compatible pair with a younger or genetically closer donor (see Roth, Sönmez, and Ünver, 2004, Ferrari et al., 2017, and Nicolò and Rodríguez-Álvarez, 2017). Such schemes not only can incentivize a limited number of compatible pairs, they can also deter participation due to uncertain and prolonged waiting times. Our proposal is the first one we are aware of that can globally and ex ante provide incentives to compatible pairs using tools that are already acceptable within the transplantation community[8].

2 A Dynamic Model of Kidney Transplantation

Consider patients who need a kidney transplant, where each patient has a blood type $X \in \{O, A, B, AB\}$. Let $\lambda_X > 0$ be the inflow rate of new type X patients; that is $\lambda_X dt$ is the measure of type X patients who enter in a small time interval dt. Suppose that the expected lifetime while living with kidney disease is distributed with a continuous and strictly increasing distribution function $F(\cdot)$ on the interval $[0, T]$. Then the measure of blood type X patients who are alive after t years is given by $\lambda_X[1 - F(t)]$. In the steady state of this model, when a transplantation option is not present, the total mass of type X patients is $\int_0^T \lambda_X[1 - F(t)]dt$.

2.1 Biological Barriers to Kidney Transplantation

The best remedy for kidney failure is transplantation. There are two potential biological barriers for kidney transplantation. A patient must be both blood-type compatible and tissue-type compatible with a potential donor to be able to receive his kidney. Type O donors are blood-type compatible with patients of all four blood-types, type A donors are blood-type compatible with patients of blood types A and AB, type B donors are blood-type compatible with patients of blood types B and AB, and type AB donors are blood-type compatible with patients of only blood type

2.2 Deceased-Donor Transplantation

The most common source of transplant kidneys in the US (and in much of the western world) is deceased donors. The United Network for Organ Sharing (UNOS) is the federal contractor in charge of allocating deceased-donor organs in the US, and it uses a point system for kidneys. Since deceased-donor organs perish within a very short period, they are allocated as soon as they are harvested. Two important features of the UNOS deceased-donor kidney allocation system are that the waiting time in the queue is the most signi cant part of the point system and kidneys are reserved for patients with the same blood type, with the exception of blood-type A kidneys that can also be allocated to blood-type AB patients.

In the medical literature, reserving blood-type X organs to blood-type X patients is referred to as ABO-identical (ABO-i) allocation policy. Since blood type AB is relatively rare, ABO-i policy is a good approximation for the allocation of deceased-donor kidneys in the US. And given the strong in uence of waiting time in the deceased-donor queue, we will assume that deceased-donor kidneys are allocated with rst-in- rst-ous6ated with)

assumption easily holds in the US and elsewhere. We assume that each patient has at most one living donor, who is of blood type X with probability $p_X > 0$, and also assume that blood types of the patient and his donor are uncorrelated. Then a blood type X patient with a living donor is (both blood-type and tissue-type) compatible with his donor with probability p^l_X, where

$$p^l_O = (1 \quad)p_O; \qquad p^l_A = (1 \quad)(p_O + p_A);$$
$$p^l_B = (1 \quad)(p_O + p_B); \text{ and } \qquad p^l_{AB} = (1 \quad)(p_O + p_A + p_B + p_{AB}) = (1 \quad):$$

We assume that a patient with a compatible living donor receives the kidney as soon as he needs a transplant without ever entering the deceased-donor queue. Then the ow of blood-type X patients who receive a living-donor transplant from their donors is

$$l_X = p^l_X \quad x \quad x:$$

Although they last longer than deceased-donor transplants, living-donor transplants can also fail. Let $^l \quad ^d$ be the steady-state fraction of living-donor transplant recipients who reenter the deceased-donor queue due to failure of their transplant. We assume that reentrants no longer have a donor upon reentry.

For each blood type X, the availability of living-donor transplantation decreases the ow of new patients to the deceased-donor queue by l_X, but a fraction of that gure, $^l l_X$, reenter due to failure of living-donor transplants. Therefore, the net steady-state ow of patients entering or reentering the blood-type X deceased-donor queue is given as

$$^{l;dec}_X = \quad _X + \quad ^d \quad _X + \quad ^l l_X \quad l_X = \quad _X + \quad ^d \quad _X \quad (1 \quad ^l)p^l_X \quad x \quad x:$$

Observe that, for each blood type X, the availability of living-donor transplantation reduces the steady-state ow of patients entering the deceased-donor queue by $(1 \ ^l)p^l_X \quad x \quad x$. Hence, the availability of living-donor transplantation uniformly bene ts all patient groups. However, when the fraction of patients with living donors is the same for all blood types, the bene t is largest for type AB and smallest for type O patients. Furthermore, since $p_A > p_B$, the bene t is larger for type A patients than for type B patients.

At any time the longest-waiting cohort of type X patients without compatible donors receive the incoming type X deceased-donor kidneys. Let this cohort have arrived $t^{l;dec}_X$ years before the current time. At steady state, we have $^{l;dec}_X[1 \quad F(t^{l;dec}_X)] = \quad _X$, and therefore the time spent on the type X deceased-donor queue by the receiving cohort can be found as

$$t^{l;dec}_X = F^{\ 1}\left(1 \quad \frac{X}{^{l;dec}_X}\right) = F^{\ 1}\left(1 \quad \frac{X}{_X + \quad ^d \quad _X \quad (1 \quad ^l)p^l_X \quad x \quad x}\right) < t^{d;dec}_X:$$

3 Kidney Exchange

While the availability of living donation bene ts all patient groups, not all willing living donors are able to donate due to an incompatibility with their intended recipients. Despite this di culty, an increasing number of patients with incompatible living donors are receiving kidney transplants through an exchange of donors with other incompatible patient-donor pairs.

Formally, a two-way kidney exchange matches two \mutually compatible" patient-donor pairs:

the patient of the first pair is compatible with the donor of the second pair, and the patient of the second pair is compatible with the donor of the first pair. Through an exchange of donors, both patients receive a kidney transplant. While patients with compatible donors can also participate in such exchanges, their participation so far has been very limited since they can directly receive a transplant from their own donors. Hence, in this section we will restrict our attention to kidney exchanges between incompatible pairs.

We consider a kidney-exchange program that operates in parallel with the deceased-donor queue. A patient with a compatible donor immediately receives a transplant from his donor without entering either the deceased-donor queue or the kidney-exchange pool. A patient with an incompatible donor, on the other hand, joins both the deceased-donor queue and the kidney-exchange pool. The patient accepts the first available kidney through these two programs.

We refer to a pair with a blood-type X patient and a blood-type Y donor as a type X-Y pair. In real-life applications, there are far fewer type A-O pairs in kidney-exchange pools than their reciprocal type, O-A pairs. The reason is that pairs of the former type are blood-type compatible, so they only join the pool when they are tissue-type incompatible. This is a relatively rare event with 0:05. Pairs of the latter type, on the other hand, are blood-type incompatible, and they always join the pool. This motivates:

Assumption 1 *For any two distinct blood types X, Y with X . Y,* $p_{X\ Y\ Y} \quad p_{Y\ X\ X}$.

That is, the in ow of type X-Y pairs (who always join the pool) is at least as much as the in ow of type Y-X pairs (who only join the pool when they are tissue-type incompatible). Since is small, this assumption easily holds in practical applications.[11]

To simplify the presentation of our analytical results, we also assume that the in ow of type B-A pairs is at least as much as the in ow of type A-B pairs. This assumption is super uous and symmetric results hold if the inequality is reversed.

Assumption 2 $p_{A\ B\ B} \quad p_{B\ A\ A}$.

Since there are far fewer type A-O pairs in the pool than their reciprocal type O-A pairs by Assumption 1, it is possible to match every A-O pair as soon as they arrive. While an arriving type A-O pair is tissue-type incompatible with a fraction of type O-A pairs in the pool, it is mutually compatible with a much larger fraction (1). Similarly, for any two distinct blood types X, Y with X . Y, it is possible to match every type Y-X pair as soon as they arrive. This is also the

instance. Moreover, this policy maximizes the mass of pairs who arrive in any interval that can be matched within that interval.

Observe that the structure of the above described optimal exchange can also accommodate FIFO matching where, whenever possible, an arriving type X-Y pair is matched with the longest waiting pair of its reciprocal type Y-X. This is the kidney-exchange mechanism we consider in this paper.

3.1 E ect of Kidney Exchange on Patients with Living Donors

The following grouping of transplants from kidney exchange is not only helpful to explain the e ect of exchange on patients with living donors, but also to highlight the welfare loss that results from excluding compatible pairs from exchange.

1. For each blood type X, transplants due to tissue-type incompatible pairs of type X-X: While these patients are blood-type compatible with their donors, they are tissue-type incompatible. Kidney exchange renders tissue-type incompatibility immaterial for them, since each one can be matched with a mutually compatible pair of identical type as soon as they join the kidney-exchange pool. The net increase in the ow of transplants at steady state is $p_{X\ X}\ _X$ for each blood type X.
2. For each pair of distinct blood types X, Y with X . Y, transplants due to tissue-type incompatible pairs of type Y-X: Tissue-type incompatibility becomes immaterial for such patients as well, since they too can be matched with a mutually compatible pair as soon as they join the kidney-exchange pool. However, the net increase in the ow of transplants at steady state is $2p_{X\ Y}\ _Y$, since each tissue-type incompatible pair of type Y-X also facilitates a transplant for a patient of its reciprocal type X-Y.
3. Transplants due to pairs of types A-B and B-A: For patients of type A-B (which has a lower in ow than type B-A under Assumption 2), both blood-type and tissue-type incompatibility become immaterial; they can immediately be matched with a pair of type B-A. The net increase in the ow of transplants at steady state is $2p_{B\ A}\ _A$, since each pair of type A-B also facilitates a transplant for a patient of type B-A.

Intuitively, kidney exchange eliminates tissue-type incompatibility from being a barrier to a living-donor transplantation, and, in doing so, it also facilitates an additional transplant to a patient with a blood-type incompatible donor. In addition, it also facilitates transplants to all patients of type A-B, and as many transplants to patients of type B-A. For pairs with O or AB patients, kidney exchange is directly tied to tissue-type incompatibility. Pairs with blood type AB patients in the kidney-exchange pool join the pool only because they are tissue-type incompatible with their own donor. Pairs with blood type O patients in the pool, on the other hand, can only receive a transplant if a mutually compatible pair of their reciprocal type becomes available for exchange through a tissue-type incompatibility. As a result, the e ect of kidney exchange on patient groups

Let e_X denote the steady-state flow of type X patients who receive a transplant through kidney exchange. For blood type O and any blood type Y, a flow $p_{O\ Y}\ _Y$ of type O-Y pairs are matched with type Y-O pairs. Therefore, $e_O = p_O(\ _O\ _O + \ _A\ _A + \ _B\ _B + \ _{AB}\ _{AB})$. Similarly,

$$e_A = p_A(\ _A\ _A + \ _{AB}\ _{AB}) + p_O\ _A\ _A + p_B\ _A\ _A;$$
$$e_B = p_B(\ _B\ _B + \ _{AB}\ _{AB}) + p_O\ _B\ _B + p_B\ _A\ _A; \text{ and}$$
$$e_{AB} = (p_{AB} + p_A + p_B + p_O)\ _{AB}\ _{AB} = \ _{AB}\ _{AB}.$$

We can summarize(O)]TJ/mt5y1.79-6 Td [&B TJ/kidneyhange5y1.79on5y1.79pairs living5y1.79donorsws:

1. Type A-B & each type X-Y Y . X:h patient (these)-306(t)27(yp)-27(es)-306(either)-306(imm)1(edia
   In either case,(O)]TJ/mty from
2. Type B-A & each type X-Y with X 6 Y and X .

For the other blood types, the e ect of kidney exchange on the deceased-donor queue is more

welfare impact. To see how the exclusion of compatible pairs a ects the gains from exchange, it is helpful to focus on the grouping in Section 3.1.

The critical types of exchanges are those in group 2: For any two distinct blood types X, Y with X . Y, a tissue-type incompatible Y-X pair exchanges its donor with a X-Y pair. To simplify the discussion, let X=O and Y=A. Through this exchange, the patient of the tissue-type incompatible A-O pair immediately receives a transplant from the donor of the O-A pair. Hence, in the presence of kidney exchange, whether the A-O pair is tissue-type incompatible or not does not a ect when or if its patient receives a transplant. But more essentially, this exchange also bene ts an A-O pair. In a way, kidney exchange transforms the \misfortune" of the A-O pair (caused by tissue-type incompatibility) to a life-saving opportunity for the O-A pair. Since the A-O pair is blood-type compatible, they would not have participated in exchange in the absence of tissue-type incompatibility. Kidney exchange not only eliminated tissue-type incompatibility as an obstacle for the transplantation, but it also facilitated a transplant for an additional patient. Put di erently, from a social-welfare point of view, there is a welfare loss when a blood-type A patient receives a transplant from a blood-type O donor. This welfare loss is avoided due to the combined roles of tissue-type incompatibility together with kidney exchange, and a more e cient utilization of living donors is obtained. But why depend on tissue-type incompatibility to avoid this welfare loss? Any pair of type A-O, whether they are tissue-type incompatible or not, can participate in kidney exchange, facilitating a transplant for an additional patient. The challenge here is that a compatible A-O pair has no reason to participate in exchange.

As our main contribution, we propose incentivizing such compatible pairs to participate in exchange by giving their patient an \insurance" against a potential future failure of his transplant. The insurance takes the form of prioritizing the patient in the deceased-donor queue in the event of a repeat kidney failure. To incentivize their participation in kidney exchange, these *prioritized reentrants* are placed at the top of the deceased-donor queue of their own blood type altering its FIFO structure. Since the welfare gains are due to inclusion of tissue-type compatible pairs of any type Y-X where Y 6= X and X . Y, we propose that the incentive scheme be provided for these pairs only. For each such pair of type Y-X, let $_{Y\ X}$ be the fraction of compatible pairs who are willing to take up the *incentivized exchange* option.

In the absence of incentivized exchange, there is an abundance of O-A pairs compared to A-O pairs. With incentivized exchange and for high values of$_{A\ O}$, this may change. We assume that compatible pairs only take the incentivized-exchange option if they can immediately participate in exchange, assuring that type A-O remains \overdemanded." We use a terminology where the parameter $_{A\ O}$ already takes this potential adjustment into consideration. Hence, we assume:

Assumption 3 *For any two distinct blood types X, Y with X . Y,*

$$[\ _{Y\ X}(1\quad) + \ ]p_{X\ Y\ Y}\quad p_{Y\ X\ X}:$$

As in the case of kidney exchange, this assumption assures that it is possible to match every Y-X pair at steady state as soon as they arrive for any two distinct blood types X, Y with X Y. And moreover, replacing Assumption 3 with Assumption 1 assures that the optimality result of Theorem

1 continues to hold under incentivized exchange. Hence, we again consider the optimal exchange mechanism where an arriving type X-Y pair, whenever possible, is matched with the longest-waiting pair of its reciprocal type, Y-X.

4.1 Effect of Incentivized Exchange on Patients with Living Donors

Since incentivized exchange simply increases the scope of kidney exchange, the analysis in this section parallels the analysis in Section 3.1. Recall that our target group for incentivized exchange is tissue-type compatible pairs of types A-O, B-O, AB-O, AB-A, and AB-B. Consider such a pair of type Y-X that takes the incentivized-exchange option. The patient of this pair could have received a transplant from his own donor, and, hence, his own transplant does not directly increase the total number of transplants. The increase is due to the patient of the reciprocal type X-Y, with whom they engage in exchange. Therefore, at steady state, the number of living transplants increases by one for each incentivized pair in the target group.

Let Y-X be any type targeted for incentivized exchange. The flow of all Y-X pairs is p_{XY}, the flow of tissue-type compatible Y-X pairs is $(1-\theta)p_{XY}$, and the flow of Y-X pairs who take the incentivized-exchange option is $\pi_{YX}(1-\theta)p_{XY}$.

For each blood type X, let i_X denote the steady-state flow of the contribution of incentivized exchange on blood-type X living-donor transplants. Living-donor transplants to blood-type A patients with AB donors increases due to incentivized pairs of type AB-A, living-donor transplants to type B patients with AB donors increases due to incentivized pairs of type AB-B, and living-donor transplants to type O patients with blood-type incompatible donors increases due to incentivized pairs of types A-O, B-O, and AB-O. Each type AB patient with a living donor already receives a living-donor transplant under kidney exchange, and, hence, living-donor transplants to type AB patients do not change. Therefore $i_{AB} = 0$,

$$i_A = \pi_{AB\,A}(1-\theta)p_{A\,AB\,AB}; \qquad i_B = \pi_{AB\,B}(1-\theta)p_{B\,AB\,AB}; \text{ and}$$
$$i_O = \pi_{A\,O}(1-\theta)p_{O\,A\,A} + \pi_{B\,O}(1-\theta)p_{O\,B\,B} + \pi_{AB\,O}(1-\theta)p_{O\,AB\,AB}.$$

Since blood type AB is rare, the flow is modest for types AB-A and AB-B. Therefore, while incentivized exchange increases living-donor transplants to patients of blood types A, B, and O, the main increase is for type O patients. We conclude this section by summarizing the effect of incentivized exchange for each patient-donor type.

1. Type A-B and each type X-Y with $Y \succ X$: Patients of these types continue to immediately receive a living-donor transplant under incentivized exchange. For those types targeted for incentivized exchange, patients of pairs who have taken this option also receive a priority increase in the deceased-donor queue for a potential future failure of their transplants.
2. Each type X-Y with $X \neq Y$ and $X \not\succ Y$: Patients of these types are the primary (but not the only) beneficiaries of incentivized exchange. For any of these types, waiting time for transplantation decreases while the fraction of its patients who receive a living-donor transplant increases. Patients of any of these types X-Y continue to join both the kidney-exchange pool and the deceased-donor queue. However, due to the increased flow of pairs from their reciprocal type

Y-X, the kidney-exchange option becomes more attractive than before with the inclusion of incentivized exchange. Therefore, patients of type X-Y are more likely to exclusively receive a transplant through kidney exchange, and thus less likely to be pooled with unpaired patients at the type X deceased-donor queue.

3. Type B-A: Patients of type B-A continue to join both the kidney-exchange pool and the type B deceased-donor queue. Incentivized exchange does not a ect the ow of the patients of type

detrimental for some: Nonprioritized & unpaired type AB patients are made worse o , and non-prioritized & unpaired type A or type B patients could be made worse o with the introduction of incentivized exchange. We next introduce a variant of incentivized exchange that does not hurt any patient population.

Consider a compatible type AB-O pair who joins the kidney-exchange pool to bene t from the insurance provided by incentivized exchange. The pair exchanges donors with a type O-AB pair, thus directly helping a type O patient. If, in the future, the incentivized patient needs a repeat transplant due to a subsequent kidney failure, he is given priority in the type AB deceased-donor queue. So participation of the AB-O pair in the pool bene ts the type O patient population but incurs a cost on the type AB patient population. Thus, what they bring to the system and what they may take from it are not \balanced." They bring an extra type O living-donor kidney to the system today, but they may take a type AB deceased-donor kidney in the future. If this feature is considered undesirable, there is an easy adjustment. Type AB patient can instead receive priority in the blood type O deceased-donor queue. And in general, the patient of any pair who participates in incentivized exchange can be given priority for the deceased-donor queue of his donor's blood type rather than his own blood type. While this modi cation changes the ABO-i feature of the deceased-donor queue, it is medically feasible since the pair is blood-type compatible. We refer to this variant of our proposed mechanism as*balanced incentivized exchange*. As we show in the next section, all patient groups weakly bene t from balanced incentivized exchange.

4.4 Equity Implications of Incentivized Exchange

We next present two results which formulate how access to living-donor and deceased-donor transplantation di er across blood types with the introduction of each transplantation modality. For these analytical results, we consider a baseline scenario where no blood type has an advantage over another for access to transplantation beyond the asymmetry induced by blood-type compatibility and the impact of the transplantation modalities analyzed.

The following result formulates how access to living-donor transplantation di ers with the successive introduction of living-donor transplantation, kidney exchange, and either version of incentivized exchange.

Theorem 2 *Suppose Assumptions 2 and 3 hold. Let* $p_A > p_B$*. Suppose that* $_X =$ *for any blood type X, and* $\frac{}{} = \frac{p_X}{p_Y}$ *for any two blood types X and Y. Suppose also that the fraction of pairs taking the incentivized exchange is uniform at a xed* < 1 *for any eligible type. Then:*

1. *For living donation only, the access to living donation is ranked as*
$$\frac{l_O}{O} < \frac{l_B}{B} < \frac{l_A}{A} < \frac{l_{AB}}{AB}:$$
2. *Kidney exchange by itself increases access to living-donor transplantation for patients of type B the most, patients of type A the next, and patients of types AB and O equally and last:* $\frac{e_B}{B} > \frac{e_A}{A} > \frac{e_{AB}}{AB} = \frac{e_O}{O}$*. With the inclusion of kidney exchange, overall access to living donation is ranked as*
$$\frac{l_O + e_O}{O} < \frac{l_B + e_B}{B} = \frac{l_A + e_A}{A} < \frac{l_{AB} + e_{AB}}{AB} = :$$

3. *Incentivized exchange (or its balanced version) by itself increases access to living-donor transplantation for patients of type O the most, patients of types A and B equally and next, and does not increase access for patients of type AB:* $\frac{i_O}{O} > \frac{i_A}{A} = \frac{i_B}{B} > \frac{i_{AB}}{AB} = 0$. *With the inclusion of either version of incentivized exchange, overall access to living donation is ranked as*

$$\frac{l_O + e_O + i_O}{O} < \frac{l_B + e_B + i_B}{B} = \frac{l_A + e_A + i_B}{A} < \frac{l_{AB} + e_{AB} + i_{AB}}{AB} = \quad :$$

Our last result formulates how access to deceased-donor transplantation di ers with the successive introduction of deceased-donor transplantation, living-donor transplantation, kidney exchange, and either version of incentivized exchange.

Theorem 3 *Suppose Assumptions 2 and 3 hold. Let* $p_A > p_B$. *Suppose that* $_X =$ *for any blood type X, and* $\frac{x}{y} = \frac{x}{y} = \frac{p_X}{p_Y}$ *for any two blood types X and Y. Suppose also that the fraction of pairs taking the incentivized exchange is uniform at a xed* < 1 *for any eligible type. Then:*

1. *With deceased-donor transplantation only, the waiting time at each deceased-donor queue is the same for any blood type X:*

$$t_O^{d;dec} = t_A^{d;dec} = t_B^{d;dec} = t_{AB}^{d;dec}.$$

2. *Introduction of living-donor transplantation reduces the waiting time at each deceased-donor queue. The changes in waiting times and the waiting times are ranked as follows:*

$$(t_{AB}^{d;dec} \quad t_{AB}^{l;dec}) > (t_A^{d;dec} \quad t_A^{l;dec}) > (t_B^{d;dec} \quad t_B^{l;dec}) > (t_O^{d;dec} \quad t_O^{l;dec});$$

$$t_{max}^{l;dec} = t_O^{l;dec} > t_B^{l;dec} (t_{AB}^{d;dec} t_A^{l;dec} > t_{AB}^{l;dec} = t_{min}^{l;dec}.$$

Further suppose that $\quad$ *and* ^{l} *are su ciently small. Then:*

$$+ \quad) > (t_B^{dec} + {}_A^{dec}$$

3. *Introduction of kidney exchange in addition to living-donor transplantation further reduces the waiting time at each deceased-donor queue, but more for blood type B than blood type A equalizing the deceased-donor queue waiting times for these two blood types. The combination of kidney exchange and living-donor transplantation reduces the waiting time at the type AB deceased-*

$$(t_{AB}^{d;dec}$$

O deceased-donor queues, while keeping it the same at the type AB deceased-donor queue. While the type A and type B deceased-donor queues continue to have equal waits, the type O deceased-donor queue continues to have the longest wait, whereas the type AB deceased-donor queue continues to have the shortest wait among the four deceased-donor queues.

The di erence between the longest and the shortest wait times decreases with the introduction of balanced incentivized exchange:

$$t_O^{b;dec} < t_O^{e;dec}; \qquad t_A^{b;dec} = t_B^{b;dec} < t_A^{e;dec} = t_B^{e;dec}; \qquad t_{AB}^{b;dec} = t_{AB}^{e;dec};$$

$$\left(t_{max}^{b;dec} = t_O^{b;dec} \right) \qquad t_{min}^{b;dec}$$

(

Model Outcomes: Patients Receiving Transplant						
	O	A	B	AB	Overall
Treatments	Living-Donor Transplants				
Living-donor transplantation (l_X)	2749.17 *18.71%*	2325.30 *23.29%*	785.76 *17.59%*	235.93 *20.30%*	6096.17 *20.12%*
Regular exchange ($e_X + l_X$)	2984.82 *20.31%*	2813.68 *28.18%*	1194.71 *26.75%*	247.65 *21.31%*	7240.85 *23.89%*
Incentivized ($e_X + l$	= 25% 3483.52 *23.71%*	2835.97 *28.41%*	1202.14 *26.92%*	247.65 *21.31%*	7769.28 *25.64%*

Model Outcomes: Average Time to Transplant for Blood-Type-Incompatible Pair Types								
Treatments		O-A	O-B	O-AB	A-B	A-AB	B-A	B-AB
Dec.&Living-Donor Trans.		pooled w O	pooled w O	pooled w O	pooled w A	pooled w A	pooled w B	pooled w B
Regular exchange		pooled w O	pooled w O	pooled w O	0	pooled w A	1.89	pooled w B
Incentivized	= 25%	pooled w O	pooled w O	pooled w O	0	pooled w A	1.89	pooled w B
(with ABO-i	= 50%	pooled w O	pooled w O/4.78	pooled w O	0	pooled w A	1.89	pooled w B
dec. donor a.)	= 100%	3.14	0.90	pooled w O /4.50	0	1.00	1.89	3.24
Incentivized	= 25%	pooled w O	pooled w O	pooled w O	0	pooled w A	1.89	pooled w B
(with de-facto	= 50%	pooled w O	4.78	pooled w O	0	4.88	1.89	pooled w B
dec. donor a.)	= 100%	3.14	0.90	4.50	0	1.00	1.89	3.24

Table 4: Model outcomes for time to transplant for blood-type-incompatible pair types (measured in years). \Pooled w X" means type X-Y pairs are pooled with type X nonprioritized & unpaired patients. This is true for both incentivized and balanced-incentivized treatments except two cases: Pairs of type O-AB are pooled with nonprioritized & unpaired O patients under ABO-i deceased-donor allocation for the incentivized treatment with = 100%, while this is not true for the balanced-incentivized treatment. The same is true for type O-B when = 50%. This happens as blood-type O nonprioritized deceased-donation recipients wait shorter times in the incentivized treatments. These are denoted in the table in those two cells with a \" sign. The pooled groups' waiting times are reported in Table 3.

Model Outcomes: Average Time to Any Type of Transplant															
Dec.	O	A	B	AB	Overall	O	A	B	AB	Overall	O	A	B	AB	Overall
Donor A.	Deceased-donor transplantation					Incentivized = 25%					Balanced inc. = 25%				
ABO-i	6.64	5.83	7.82	7.90	6.51	3.40	2.69	3.49	3.85	3.16	3.41	2.67	3.50	3.88	3.16
De facto	6.93	5.98	7.28	4.79	6.51	3.51	2.74	3.36	2.72	3.17	3.52	2.72	3.36	2.68	3.17
	Living-donor transplantation					Incentivized = 50%					Balanced inc. = 50%				
ABO-i	3.75	3.02	4.29	3.99	3.55	3.19	2.71	3.50	3.82	3.07	3.22	2.68	3.51	3.88	3.08
De facto	3.86	3.08	4.11	2.75	3.55	3.30	2.72	3.37	2.75	3.08	3.32	2.73	3.37	2.68	3.09
	Regular exchange					Incentivized = 100%					Balanced inc. = 100%				
ABO-i	3.61	2.66	3.49	3.88	3.24	2.71	2.66	3.41	3.72	2.81	2.78	2.61	3.45	3.88	2.84
De facto	3.72	2.71	3.35	2.68	3.24	2.81	2.71	3.30	2.81	2.84	2.86	2.66	3.31	2.68	2.84

Table 5: Model outcomes for average waiting time for any type of transplant for di erent patient blood types (measured in years).

We summarize our main ndings regarding the equity consequences of di erent transplantation policies as follows:

De-facto deceased-donor and living-donor transplantation together help type AB patients the most, followed by type A and then by type O and type B. While 67.98 percent of all AB patients and 61.51 percent of all A patients bene t from these two modalities, less than 48 percent of all B patients and 51 percent of all O patients receive transplant (these are the fractions $\frac{{}^{0}x + I_x}{x}$; see the \Total Transplants" section of Table 2).

At the margin, type B patients bene t the most from regular exchange. Type A patients bene t next most, while type O and type AB patients bene t the least. While 9.15 percent more of

The phrase \altruistically unbalanced" re ects their dismissal of the concept. Their concern is potential coercion of compatible pairs who have nothing to gain from exchange. In contrast, exchange is no longer \altruistically unbalanced" under incentivized exchange, since patients of participating pairs are insured against a repeat failure.

Under an *indirect exchange*, the donor of an incompatible pair donates a kidney to the deceased-donor queue in exchange for a priority for his patient in the deceased-donor queue. Hence this variant involves an exchange between an incompatible pair and the deceased-donor queue. Ross and Woodle (2000) object to indirect exchange for blood-type incompatible pairs, but support it for blood-type compatible (but tissue-type incompatible) pairs. Consider a type O patient with a type A donor. Under an indirect exchange, the pair donates a type A kidney to the donor queue in exchange for priority for a type O deceased-donor kidney. That is, they receive priority for a more

queue in case the patient cannot be placed at the end of a kidney chain. Conceptually incentivized exchange is similar, but it evades the above-mentioned shortcomings since incentivized pairs are blood-type compatible.

In summary, incentivized exchange harbors all the positive elements of the above variants of kidney exchange without su ering from their shortcomings.

References

Agarwal, Nikhil, Itai Ashlagi, Eduardo Azevedo, Clayton Featherstone, and Omer Karaduman (2017). \Market failure in kidney exchange." Working paper.

Akbarpour, Mohammad, Shengwu Li, and Shayan Oveis-Gharan (2017). \Thickness and information in dynamic matching markets." Working paper.

Anderson, Ross, Itai Ashlagi, David Gamarnik, and Yash Kanoria (2017). \E cient dynamic barter exchange."*Operations Research*, forthcoming.

Lee, SangMok (2017). \Incentive compatibility of large centralized matching markets."*Review of Economic Studies*, 84 (1), 444{463.

Liu, Qingmin and Marek Pycia (2013). \Ordinal e ciency, fairness, and incentives in large markets." Working paper.

Nicob, Antonio and Carmelo Rodrguez-Alvarez (2017). \Age-based preferences in paired kidney exchange."*Games and Economic Behavior*, 102, 508{524.

Rapaport, F. T. (1986). \The case for a living emotionally related international kidney donor exchange registry."*Transplantation Proceedings*, 18, 5{9.

Ratner, Lloyd E., Abbas Rana, Emily R. Ratner, et al. (2010). \The altruistic unbalanced paired kidney exchange: proof of concept and survey of potential donor and recipient attitudes."*Transplantation*

For Online Publication

Appendix A Omitted Proofs and Additional Results on Waiting Times

In this appendix, we provide the omitted proofs and some additional results on waiting times. To do that, we rst provide a formal model of tissue-type incompatibility: Each patient has a type, depending on against which subset of HLA tissue proteins he has preformed antibodies. We study the limit as the number of types goes to in nity. First, x the number of types to a y

with any type pair. A self-demanded-type pair can be matched with a same-type or overdemanded-type pair. In particular, the following results hold:

A.2 Pooling and Waiting Times under Regular and Incentivized Exchange

Recall that for all incentivized-exchange-eligible pairs, i.e., of all types X-Y such that Y $\in$ X, Y $\notin$ X, and the patient and donor have no tissue-type incompatibility, $\alpha_{XY} \in [0,1]$ is the fraction that participate in incentivized exchange. Let $\alpha = (\alpha_{XY})_{Y \in X; Y \notin X}$ be the vector of such fractions. We use the terms "regular exchange" and "incentivized exchange with $\alpha = 0\%$" interchangeably. To determine the steady-state outcomes, we introduce certain flow rates.[16] For each blood type X and each Y $\in$ X, let

$$i_{XY} = \begin{cases} [\lambda_{XY} + \alpha_{XY}(1-\theta)]p_Y \, r_X \, r_Y & \text{if } Y \in X \\ p_Y \, r_X \, r_Y & \text{otherwise} \end{cases} \tag{1}$$

refer to the pair-type X-Y flow to the exchange pool. Let the *incentivized pair flow* relevant for blood type X be given by

$$c_X = \sum_{Y : Y \in X \,\&\, Y \notin X} \alpha_{XY}(1-\theta)p_Y \, r_X \, r_Y. \tag{2}$$

Observe that $^{I}c_X$ is the reentry flow of previously incentivized type X patients. These patients will be prioritized in the deceased-donor queue of blood type X and will not wait upon reentry. Thus, the effective flow rate of deceased-donor kidneys for nonprioritized type X patients is $s_X - {}^{I}c_X$. We also have

$$i_X = \underbrace{(1-\alpha_X)\lambda_X}_{\text{new unpaired}} + \underbrace{d_X}_{\text{reentry / deceased}} + \underbrace{{}^{I}[l_X + e_X + i_X - c_X]}_{\text{reentry / all live minus incentivized}} \tag{3}$$

as the total *nonprioritized & unpaired type X patient flow*.

We define the following ratios:

1. The ratio of the deceased-donor effective flow for nonprioritized patients to the nonprioritized & unpaired patient flow is

$$r_X = \frac{s_X - {}^{I}c_X}{i_X}. \tag{4}$$

2. For each underdemanded type X-Y except B-A (i.e., X $\in$ Y and X $\notin$ Y), the ratio of the flow of incompatible or incentivized type Y-X pairs to the total flow of type X-Y pairs is

$$r_{XY} = \frac{i_{YX}}{i_X} = \frac{[\lambda_{YX} + \alpha_{YX}(1-\theta)]p_Y \, r_Y \, r_X}{p_X \, r_X \, r_Y} = \frac{[\lambda_{YX} + \alpha_{YX}(1-\theta)]p_Y}{p_X}$$

9701 Tf 5.875 -1.793 Td [0 Td [( )]TAf 5.59 0 Td [(Y)]TJ/F58d [(=)]TJ/F20 111.9552 Tf 7.14 -13.
-7.359 Td [(Y)]TJ/F51 7.9701 Tf 6.0
9701 Tf 5.875 -1.793 Td [( )]TJ/F20 11.9552 Tf 7.288 15.064 Td [( )]TJ/F20 11.9552 Tf 7.14 -13.27

for nonprioritized patients to nonprioritized & unpaired type X patients. For an underdemanded type X-Y, ratio $r_{X\ Y}$ would be relevant if type X-Y pairs did not receive any deceased donation but only participated in exchange. In these cases, the waiting time for a deceased-donor transplant for nonprioritized & unpaired type X patients would be

$$t_X = F^{\ 1}\ 1 \qquad {}^{X} \qquad {}^{I}c_X$$

1. Blood-type X patients, who are in overdemanded-type or self-demanded-type pairs and who have either incompatible donors or are eligible and willing to participate in incentivized exchange, participate in exchange immediately upon their arrival to the exchange pool.
2. Suppose the patients in pairs of underdemanded types X-Y_1, ..., X-$Y_{\grave{}(X)}$ receive both deceased-donor and exchange transplants, while the patients in pairs of underdemanded types X-$Y_{\grave{}(X)+1}$, ..., X-$Y_{k(X)}$ receive only exchange transplants for some $\grave{}(X) \in \{0, \ldots, k(X)\}$. Then nonprioritized & unpaired type X patients and the patients of type X-Y_1, ..., X-$Y_{\grave{}(X)}$ pairs wait for a deceased-donor (or exchange) transplant for the duration

$$t_X^{i;dec} = F^{-1}\left(1 - \frac{{}^{X}c_X + {}^{i}_{Y_1}{}_{X} + \cdots + {}^{i}_{Y_{\grave{}(X)}}{}_{X}}{{}^{i}_{X} + {}^{i}_{X}{}_{Y_1} + \cdots + {}^{i}_{X}{}_{Y_{\grave{}(X)}}}\right); \text{ and,} \quad (6)$$

for all $\grave{}$

thus, $r_{A;A\ AB\ =0} = r_{B;B\ AB\ =0}$ implying that

$$t_A^{e;dec} = F^{\ 1}\ 1\quad r_{A;A\ AB\ =0} = F^{\ 1}\ 1\quad r_{B;B\ AB\ =0} = t_B^{e;dec}:$$

Suppose $^{\ l} = 0$. Then,

$$r_{O;O\ A;O\ B;O\ AB\ =0;\ ^{\ l}=0} = \frac{\frac{\ O}{O} + (\ p_A\ +\ p_B\ +\ p_{AB}\ )}{1\quad +\quad d\frac{\ O}{O} + (p_A\ +\ p_B\ +\ p_{AB}\ )};$$

$$r_{A;A\ AB\ =0;\ ^{\ l}=0} = \frac{\frac{\ A}{A} + (\ p_{AB}\ )}{1\quad +\quad d\frac{\ A}{A} + (p_{AB}\ )}; \text{ and}$$

$$r_{AB\ \ ^{\ l}=0} = \frac{\frac{\ AB}{AB}}{1\quad +\quad d\frac{\ AB}{AB}}:$$

Since for su ciently small $\ ,\ \frac{x=\ x}{1\quad +\ d\ x=\ x} >\ $ for any X, we have that $\frac{x=\ x\ +\ q(X)}{1\quad +\ d\ x=\ x\ +q(X)}$ is decreasing in $q(X)$ for any $q(X)\quad 0$. Thus, we can rank the above entities as $r_{O;O\ A;O\ B;O\ AB\ =0;\ ^{\ l}=0} < r_{A;A\ AB\ =0;\ ^{\ l}=0} < r_{AB\ \ ^{\ l}=0}$. By the continuity of these r ratios in $^{\ l}$, for su ciently small $^{\ l}$ we still have $r_{O;O\ A;O\ B;O\ AB\ =0} < r_{A;A\ AB\ =0} < r_{AB}$. As the generic waiting time$t = F^{\ 1}(1\quad r)$ is decreasing in$r$, we can rank the waiting times for deceased donation in the queue unde p

(b) Introduction of balanced incentivized exchange, in addition to regular exchange, direct living-donor transplantation, and deceased-donor transplantation, causes the waiting time for a deceased-donor transplant to decrease for all blood types except AB, for which it remains constant by Lemma 2, with respect to regular exchange.

We next inspect the magnitude of these changes.

Note that $b_X = i_X$ for each blood type X, i.e., the same flow of pairs benefit from both balanced-incentivized-exchange and incentivized-exchange regimes. Given a blood type X, the relevant ratio for waiting times in the deceased-donor queue, which we will refer to as r_X for short, satisfies

$$r_X := r^b_{X;X\ Y_1;\dots;X\ Y_{b(X)}} = \frac{\frac{\cdot_X}{\cdot_X}\left[\frac{c^b_X}{\cdot_X} + \sum_{m=1}^{b(X)}\left([\ \ + (1\ \ )]p_{Y_m}\right)\right]}{1 + \left[\frac{d_X}{\cdot_X} + \frac{l_X + e_X + i_X}{\cdot_X} + \sum_{m=1}^{b(X)} p_{Y_m}\right]}:$$

We prove several claims:

Claim 1. Deceased-donation waiting times in the A and B deceased-donor queues are the same under incentivized balanced exchange, i.e., $t^{b;dec}_A = t^{b;dec}_B$.

Proof of Claim 1. We have $\frac{l_O + e_O + i_O}{\cdot_O} < \frac{l_A + e_A + i_A}{\cdot_A} = \frac{l_B + e_B + i_B}{\cdot_B} < \frac{l_{AB} + e_{AB} + i_B}{\cdot_{AB}} =$ by Theorem 2 and $\frac{\cdot_X}{\cdot_Y} = \frac{\cdot_X}{\cdot_Y}$ for any two blood types X and Y. Moreover, c^b_X

for each blood type O, A, and AB as follows for any l:

$$r_O = \frac{\frac{O}{O} \; l \; (1 \quad)(p_A + p_B + p_{AB}) \; + \; P_{m=1}^{\; b(O)} \quad + \; (1 \quad) \; p_{Y_m}}{1 \quad + \; d\frac{O}{O} + \; l \; p_O \; + \quad + \; (1 \quad) \; (p_A + p_B + p_{AB}) \quad + \; P_{m=1}^{\; b(O)} p_{Y_m}};$$

$$r_A = \frac{\frac{A}{A} \; l \; (1 \quad)p_{AB} \; + \; P_{m=1}^{\; b(A)} \quad + \; (1 \quad) \; p_{Y_m}}{1 \quad + \; d\frac{A}{A} + \; l \; (p_O + p_A + p_B) l \; +}$$

all underdemanded pair types X-Y except B-A (where equalities follow by Equations 19 and 17). Thus, $r_{X\ |=0}$ is decreasing in $q(X)$. Since

$$q(AB) = 0 < q(A) = p_{AB} < q(O) = (p_A + p_B + p_{AB}) ;$$

we obtain

$$r_{O\ |=0} < r_{A\ |=0} < r_{AB\ |=0}:$$

Finally, each of these is increasing in

Hence, by the Taylor series expansion of the ratios in $(\ ^{l};\ )$ around $(0;0)$, for sufficiently small $(\ ^{l};\ )$ we have $r_O < r_A < r_{AB}$. Since $t_X^{b;dec} = F^{-1}(1 - r_X)$ and by Claim 1 $t_A^{b;dec} = t_B^{b;dec}$, we obtain

$$t_O^{b;dec} < t_A^{b;dec} = t_B^{b;dec} < t_{AB}^{b;dec}.$$

Finally, we complete the proof of Part 4(b) and the theorem as follows: Since $t_{min}^{b;dec} = t_{AB}^{b;dec} = t_{AB}^{e;dec} = t_{min}^{e;dec}$ and $t_{max}^{b;dec} = t_O^{b;dec} < t_O^{e;dec} = t_{max}^{e;dec}$, we obtain

$$t_{max}^{b;dec} - t_{min}^{b;dec} < t_{max}^{e;dec} - t_{min}^{e;dec}.$$

■

Appendix B Construction of Calibration Parameters for Numerical Predictions

In this appendix, we explain how the calibration parameters, reported in Table 1 in Section 5 and used to generate the numerical model predictions, are constructed.

In Table 6, we report the blood-type distribution for different ethnicities and fractions 4 -245915254e

The US OPTN Kidney Data					
	O	A	B	AB	Overall
Total Additions to the Queue	16323	11090	4919	1325	33657
Living-Donation Recipients not on Queue	432	406	127	35	1000
Reentrants	2062	1513	580	198	4353
Direct Living-Donation Recipients	2750	2326	786	236	5098
Exchange Participants	128	96	58	8	290
Deceased-Donation Recipients	4726	3815	1347	554	10442
Deceased Donors	3458	2722	850	218	7248
Average CPRA					4.73%

Table 7: Arrivals to and transplants from the kidney deceased-donor queue for 2009 entrants. Year 2009 is used because this is the latest year for which five-year dialysis survival rates are available as of May, 2017. Data is obtained from the OPTN using the "national data" option from http://optn.transplant.hrsa.gov 05/12/2017.

transplants without even registering in the queue (the second row of the table). We add these two numbers and subtract the number of reentrants (the third row of the table) from them to find ν_X for each blood type X.

Reentry fractions, ρ^l and ρ^d, are assumed to be the same, as the OPTN national data do not distinguish reentrants based on their previous transplantation type. We divide the total number of reentrants (the last cell of the third row of the table) by the sum of direct living-donation recipients, exchange participants, and deceased-donation recipients (the sum of the last cells of the fourth, fifth, and sixth lines of the table).

The tissue-type incompatibility probability, π, is taken as the average calculated panel reactive antibody (CPRA), 0.0473, for the 2009 entrants (see Table 9 in Ap2

Y 6 X, $_{X\ Y}$, is our free calibration variable. We assume that this fraction is uniform for each type, and we denote it as . We consider three treatments with = 25%, 50%, and 100%.

US On-dialysis Survival Rates					
Time:	3 mo.	1 yr.	2 yr.	3 yr.	5 yr.
On dialysis	0.9215	0.7824	0.6648	0.5694	0.4230

8: For the survival rate, 1 F(t), we use the on-dialysis survival probability of US kidney disease patients for the 2009 entrant cohort. These 2009 estimates for dialysis patients are obtained from the weighted average of hemodialysis and peritoneal dialysis survival rates reported in the 2016 Annual Report of the National Kidney Organization, Volume 2, Table 6.3 of Chapter 6 (retrieved from https://www.usrds.org/2016/download/v2_c06_Mortality_16.pdf on 05/12/2017). The weights used in the weighted average calculations are the average 2009{2014 percentages of patients on hemodialysis versus peritoneal dialysis, reported in the same report's Volume 2, Figure 1.2 of Chapter 1 (retrieved from https://www.usrds.org/2016/download/v2_c01_IncPrev_16.pdf on 05/12/2017).

In Table 8, survival rates over time, 1 F(t), are reported. These are obtained from the US Renal Data System (USRDS) data for dialysis patients. We t an exponential duration curve (for t measured in years) as

$$\hat{F}(t) = 1 \quad \hat{a}e^{\hat{\delta}t}$$

and obtain the following estimates using non-linear least squares regression (through the command in MATLAB): $\hat{a} = 0{:}9427$ with the 95 percent con dence interval (0:8945; 0:9909), and $\hat{\delta} = 0$:

We assume patients are heterogenous in their tissue-type incompatibility probabilities. We use the entrant CPRA distribution reported in Table 9 to generate the tissue-type incompatibility probability $_i$ for each patient i. The mean of this distribution gives us the value of we use in the numerical model predictions, 0.0473 (or 473 percentage points in the CPRA reporting metric), in Section 5. This data gives the fraction of entrants in ve di erent CPRA intervals. We assume that all patients uniformly and randomly take CPRA values in their assigned CPRA intervals. For example, the US OPTN data reports that 1.73 percent of all entrants have CPRA points between 0 percent and 20 percent (the second column of this table). We rst assume that a simulated patient i is assigned to this group with probability 0.0173. Then his exact tissue-type incompatibility probability $_i$ is determined uniformly randomly from the interval $(0; 0.2)$.

The US OPTN Data for CPRA Distribution for Entrants					
	CPRA intervals (in % points)				
	0	(0,20)	[20,80)	[80,98)	[98,100)
Fraction of Entrants	91.57%	1.73%	3.85%	1.88%	0.98%

Table 9: Data is obtained from the OPTN using the "national data" option for the year 2009 from http://optn.transplant.hrsa.gov on 03/28/2018.

Our simulations use a scaled-down version of the calibrated in ow rates for new patients and deceased donors. The US consists of 13 transplant regions of various sizes. Deceased-donor kidneys are rst o ered to patients within their arrival regions. If a suitable match cannot be found in the region, then they are o ered nationally. Our simulation roughly maps to one region that comprises

average waiting time of patients who arrive in year 31. Thus, when the simulation ends at year 55, all patients

Simulation Results: Patients Receiving Transplant						
		O	A	B	AB	Overall
Treatments		Living-Donor Transplants				
Living-donor transplantation (l_X)		273.92 *18.68%* (16.00) *(0.99%)*	232.82 *23.22%* (14.15) *(1.32%)*	76.81 *17.33%* (9.11) *(1.81%)*	22.98 *19.82%* (4.93) *(3.99%)*	606.53 *20.03%* (23.02) *(0.75%)*
Regular exchange ($e_X + l_X$)		290.70 *19.83%* (17.20) *(1.08%)*	278.95 *27.82%* (16.39) *(1.50%)*	117.18 *26.46%* (11.73) *(2.39%)*	23.90 *20.61%* (5.06) *(4.09%)*	710.73 *23.48%* (28.64) *(0.93%)*
Balanced	= 25%	340.98 *23.26%* (17.12) *(1.11%)*	280.78 *28.00%* (16.99) *(1.57%)*	117.98 *26.64%* (12.00) *(2.47%)*	23.93 *20.64%* (5.02) *(4.06%)*	763.67 *25.22%* (31.59) *(1.03%)*
Incentivized ($e_X + l_X + b_X$)	= 50%	390.20 *26.62%* (18.56) *(1.24%)*	282.94 *28.22%* (16.77) *(1.55%)*	118.51 *26.76%* (12.01) *(2.47%)*	23.88 *20.59%* (5.10) *(4.10%)*	815.53 *26.94%* (33.30) *(1.08%)*
	= 100%	487.91 *33.29%* (20.71) *(1.42%)*	287.33 *28.66%* (17.08) *(1.56%)*	119.96 *27.09%* (11.93) *(2.46%)*	23.88 *20.59%* (5.07) *(4.06%)*	919.08 *30.36%* (38.03) *(1.23%)*
Balanced Incentivized also with comp. type X-X pairs ($e_X + l_X + b_X$)	= 25%	344.85 *23.53%* (17.01) *(1.11%)*	282.39 *28.16%* (16.93) *(1.55%)*	118.31 *26.71%* (11.93) *(2.46%)*	23.96 *20.66%* (5.08) *(4.10%)*	769.51 *25.42%* (31.58) *(1.03%)*
	= 50%	395.16 *26.96%* (18.17) *(1.21%)*	285.05 *28.43%* (16.93) *(1.54%)*	119.04 *26.88%* (11.91) *(2.43%)*	23.91 *20.61%* (5.10) *(4.10%)*	823.16 *27.19%* (32.52) *(1.05%)*
	= 100%	493.40 *33.67%* (20.92) *(1.43%)*	289.38 *28.86%* (17.38) *(1.59%)*	120.19 *27.14%* (11.79) *(2.44%)*	23.90 *20.61%* (5.11) *(4.12%)*	926.87 *30.61%* (38.40) *(1.24%)*
Treatments		Deceased-Donor Transplants				
Non-incentivized Treatments		474.77 *32.41%* (20.80) *(1.72%)*	380.82 *38.02%* (18.32) *(2.29%)*	134.51 *30.43%* (12.06) *(3.05%)*	57.23 *49.69%* (7.14) *(7.55%)*	
Balanced	= 25%	466.92 *31.87%* (21.13) *(1.73%)*	384.71 *38.40%* (18.63) *(2.33%)*	137.26 *31.05%* (12.19) *(3.12%)*	58.44 *50.73%* (7.18) *(7.60%)*	
	= 50%	459.08 *31.34%* (12.06) *(4.12%)*	388.21 *38.75%* (5.07) *(4.06%)*	147 11 522 0 Td [((11.91))]TJ/F63 11.9552 Tf 41.4372 0 Td [((11.91))]TJ/F63 111.211[(20638))]TJ/F63		

Simulation Results: Average Time to Nonprioritized Deceased-Donor Transplant															
O	A	B	AB	Overall	O	A	B	AB	Overall	O	A	B	AB	Overall	
Deceased-donor transplantation					Balanced inc. = 25%					Balanced inc. also with X-X = 25%					
6.893	5.949	7.235	4.766	6.474	5.491	4.689	5.711	3.799	5.131	5.490	4.686	5.720	3.823	5.132	
(0.105)	(0.119)	(0.193)	(0.350)	(0.064)	(0.101)	(0.124)	(0.189)	(0.316)	(0.066)	(0.098)	(0.125)	(0.205)	(0.331)	(0.066)	
Living-donor transplantation					Balanced inc. = 50%					Balanced inc. also with X-X = 50%					
6.074	4.928	6.459	3.897	5.585	5.107	4.678	5.667	3.768	4.947	5.111	4.674	5.667	3.758	4.948	
(0.103)	(0.126)	(0.196)	(0.288)	(0.068)	(0.103)	(0.128)	(0.175)	(0.292)	(0.065)	(0.098)	(0.125)	(0.189)	(0.306)	(0.066)	
Regular exchange					Balanced inc. = 100%					Balanced inc. also with X-X = 100%					
5.932	4.707	5.752	3.837	5.347	4.997	4.699	5.630	3.684	4.895	5.009	4.691	5.620	3.680	4.895	
(0.108)	(0.125)	(0.190)	(0.325)	(0.070)	(0.105)	(0.127)	(0.180)	(0.319)	(0.070)	(0.107)	(0.136)	(0.185)	(0.309)	(0.072)	

Table 11: Simulation results under the de-facto allocation policy for deceased-donor waiting time for nonprioritized patients from different blood types (measured in years).

Simulation Results: Average Time to Any Type of Transplant															
O	A	B	AB	Overall	O	A	B	AB	Overall	O	A	B	AB	Overall	
Deceased-donor transplantation					Balanced inc. = 25%					Balanced inc. also with X-X = 25%					
6.893	5.949	7.235	4.766	6.474	3.587	2.676	3.265	2.597	3.159	3.548	2.655	3.258	2.599	3.133	
(0.105)	(0.119)	(0.193)	(0.350)	(0.064)	(0.107)	(0.111)	(0.157)	(0.261)	(0.066)	(0.101)	(0.106)	(0.161)	(0.274)	(0.065)	
Living-donor transplantation					Balanced inc. = 50%					Balanced inc. also with X-X = 50%					
3.862	3.057	4.095	2.743	3.537	3.401	2.626	3.191	2.517	3.051	3.362	2.602	3.175	2.508	3.023	
(0.118)	(0.125)	(0.206)	(0.252)	(0.076)	(0.100)	(0.111)	(0.176)	(0.260)	(0.067)	(0.091)	(0.109)	(0.157)	(0.261)	(0.062)	
Regular exchange					Balanced inc. = 100%					Balanced inc. also with X-X = 100%					
3.766	2.720	3.331	2.654	3.258	2.942	2.522	3.057	2.386	2.787	2.912	2.504	3.032	2.383	2.762	
(0.114)	(0.110)	(0.171)	(0.280)	(0.067)	(0.097)	(0.109)	(0.168)	(0.252)	(0.064)	(0.092)	(0.109)	(0.161)	(0.242)	(0.061)	

Table 12: Simulation results under the de-facto deceased-donor allocation policy for average waiting time for any type of transplant for different patient blood types (measured in years).

Appendix D Perfect Matching with (Heterogenous) Tissue-Type Incompatibilities

Simulation Results: Average Time to Nonprioritized Deceased-Donor Transplant															
O	A	B	AB	Overall	O	A	B	AB	Overall	O	A	B	AB	Overall	
Deceased-donor transplantation					Balanced inc. = 25%					Balanced inc. also with X-X = 25%					
6.893	5.949	7.235	4.766	6.474	5.491	4.689	5.711	3.799	5.131	5.490	4.686	5.720	3.823	5.132	
(0.105)	(0.119)	(0.193)	(0.350)	(0.064)	(0.101)	(0.124)	(0.189)	(0.316)	(0.066)	(0.098)	(0.125)	(0.205)	(0.331)	(0.066)	
Living-donor transplantation					Balanced inc. = 50%					Balanced inc. also with X-X = 50%					
6.074	4.928	6.459	3.897	5.585	5.107	4.678	5.667	3.768	4.947	5.111	4.674	5.667	3.758	4.948	
(0.103)	(0.126)	(0.196)	(0.288)	(0.068)	(0.103)	(0.128)	(0.175)	(0.292)	(0.065)	(0.098)	(0.125)	(0.189)	(0.306)	(0.066)	
Regular exchange					Balanced inc. = 100%					Balanced inc. also with X-X = 100%					
5.932	4.707	5.752	3.837	5.347	4.997	4.699	5.630	3.684	4.895	5.009	4.691	5.620	3.680	4.895	
(0.108)	(0.125)	(0.190)	(0.325)	(0.070)	(0.105)	(0.127)	(0.180)	(0.319)	(0.070)	(0.107)	(0.136)	(0.185)	(0.309)	(0.072)	

Table 11: Simulation results under the de-facto allocation policy for deceased-donor waiting time for nonprioritized patients from di erent blood types (measured in years).

Simulation Results: Average Time to Any Type of Transplant															
O	A	B	AB	Overall	O	A	B	AB	Overall	O	A	B	AB	Overall	
Deceased-donor transplantation					Balanced inc. = 25%					Balanced inc. also with X-X = 25%					
6.893	5.949	7.235	4.766	6.474	3.587	2.676	3.265	2.597	3.159	3.548	2.655	3.258	2.599	3.133	
(0.105)	(0.119)	(0.193)	(0.350)	(0.064)	(0.107)	(0.111)	(0.157)	(0.261)	(0.066)	(0.101)	(0.106)	(0.161)	(0.274)	(0.065)	
Living-donor transplantation					Balanced inc. = 50%					Balanced inc. also with X-X = 50%					
3.862	3.057	4.095	2.743	3.537	3.401	2.626	3.191	2.517	3.051	3.362	2.602	3.175	2.508	3.023	
(0.118)	(0.125)	(0.206)	(0.252)	(0.076)	(0.100)	(0.111)	(0.176)	(0.260)	(0.067)	(0.091)	(0.109)	(0.157)	(0.261)	(0.062)	
Regular exchange					Balanced inc. = 100%					Balanced inc. also with X-X = 100%					
3.766	2.720	3.331	2.654	3.258	2.942	2.522	3.057	2.386	2.787	2.912	2.504	3.032	2.383	2.762	
(0.114)	(0.110)	(0.171)	(0.280)	(0.067)	(0.097)	(0.109)	(0.168)	(0.252)	(0.064)	(0.092)	(0.109)	(0.161)	(0.242)	(0.061)	

Table 12: Simulation results under the de-facto deceased-donor allocation policy for average waiting time for any type of transplant for di erent patient blood types (measured in years).

Appendix D Perfect Matching with (Heterogenous) Tissue-Type Incompatibilities

$$(1 \quad) \sum_{i=1}^{k} \frac{m_{(i);k}(\quad_{(i);k} + (1 \quad_{(i);k}))}{M} \sum_{i=1}^{k} m_{(i);k}[1 \prod_{j=1;j\neq i} (1 \quad (1 \quad_{(j);k})(1 \quad_{(i);k}))];$$

where $M = \sum_{i=1}^{k} m_{(i);k}(\quad_{(i);k} + \quad (1 \quad_{(i);k}))$.

For overdemanded-type pairs, only tissue-type incompatible ones participate in the regular exchange. However, in the incentivized exchange, compatible pairs also participate. As a result, a fraction of the overdemanded pairs are compatible, while the rest are incompatible. Here, is the participation rate of compatible pairs. The assumption guarantees that, for any set of overdemanded-type pairs, the set of underdemanded pairs that are compatible with at least one pair in the set has a greater measure as $\to 0$.

Lemma 5 *Suppose Assumption 6 holds. Consider two measurable sets of overdemanded X-Y pairs and underdemanded Y-X pairs with the same measure. Suppose that a fraction of overdemanded X-Y pairs are known to be tissue-type incompatible and the rest are known to be tissue-type compatible, but otherwise these sets are formed randomly using the governing population distributions. Then, as the number of patient types* k *goes to in nity, almost every pair can be matched with a compatible pair.*

Proof.

of type (i). On the other hand, if $i > l$, then the measure of underdemanded Y-X pairs with patient type (i) who are incompatible with all such pairs is $m_{(i);k} \prod_{j=1}^{l}(1 - (1 - _{(j);k})(1 - _{(i);k}))$. Hence, the measure of underdemanded Y-X pairs that are compatible with at least one overdemanded X-Y pair in the chosen set is at least $\sum_{i=1}^{k} m_{(i);k}[1 - \prod_{j=1;j\neq i}^{l}(1 - (1 - _{(j);k})(1 - _{(i);k}))]$. This sum is greater than the measure of chosen overdemanded X-Y pairs Assumption 6.

The proof that 1 measure of overdemanded X-Y pairs can be matched follows. The lemma follows by taking $k \to 1$ and $\to 0$. ■

D.4 Matching Self-Demanded-Type Pairs

In this section, we consider the case when we match self-demanded type pairs. Fix any self-demanded-type pair X-X for some blood type X. Any such pair in the exchange pool is tissue-type incompatible. We match these pairs with each other. Therefore, in contrast with the previous sections, this is a one-sided matching problem.

We make the following assumption to show that almost every pair can be matched in the limit.

Assumption 7 *For every* > 0*, there exists* $k_0 \in \mathbb{N}$ *such that for every* $k > k_0$, $l \leq k$*, and every permutation of patient types,*

$$(1 -) \sum_{i=1}^{l} m_{(i);k} _{(i);k} \leq \sum_{i=1}^{k} m_{(i);k} _{(i);k}[1 - \prod_{j=1;j\neq i}^{l}(1 - (1 - _{(j);k})(1 - _{(i);k}))].$$

Our next result shows that under this assumption almost all self-demanded pairs can be matched.

Lemma 6 *Suppose Assumption 7 holds. Consider a set of self-demanded-type pairs X-X that are tissue-type incompatible. Assume that this set is formed randomly using the governing population distributions. Then, as the number of patient types* k *goes to in nity, almost every pair can be matched with a compatible pair.*

Proof. Since the pairs are tissue-type incompatible, but otherwise formed randomly using the governing population distributions, for each patient type i, the measure of pairs with patient type i is proportional to $m_{i\ i}$.

Fix a small > 0. Consider any k that satis es Assumption 7.

We use Gale's Supply-Demand Theorem (Gale, 1957) to show that 1 fraction of the self-demanded X-X pairs can be matched with compatible self-demanded X-X pairs. To show this, we rst construct a two-sided matching problem with these pairs. For any patient type i, we split the set of pairs with patient type i into two sets with equal measure. These sets are then added to di erent sides of the market. As a result, we get a two-sided matching problem where each side has X-X pairs where those with patient type i have a measure proportional to $m_{i\ i}$. For ease of exposition, suppose that the measure is exactly $m_{i\ i}$.

Consider one side of the market. To apply Gale's Supply-Demand Theorem, take a random measurable subset of pairs on this side of the market that has measure 1 fraction of all pairs on this side. Since the subset is chosen randomly, the compatibility of patients can still be formed randomly using the governing population. We need to show that for any subset of pairs, the measure

of pairs on the other side of the market that are compatible with at least one pair in the set is weakly

whereas the right side is positive. Furthermore, as $k \to 1$ Inequality 23 holds also for every $j \leq n$ because $m_{i;k} \to 0$ for every i and n is a fixed natural number which does not depend on k. In this case, the left side converges to zero and the right side is always positive. ■